UDC 633.2: 633.3

PRODUCTIVITY OF LONG-TERM BEAN GRASS IN SINGLE-SPECIES AND COMPATIBLE WITH ANWLESS BROMEGRASS FEED AGROPHYTOCENOSES

V.H. Kurhak, S.M. Panasyuk,

NSC «Institute of Agriculture NAAS» (Chabany, Ukraine)

Goal. Establish the best perennial legumes when grown in single-species and compatible fodder agroecosystems with boneless and the effectiveness of the use of nitrogen-fixing and phosphorus-mobilizing bacterial preparations, as well as liming. **Methods.** General scientific – hypotheses, inductions and deductions, analogies, generalizations and special – field, laboratory, mathematical and statistical, computational and comparative. **Results.** Based on the generalization of literature sources, it is proved that the creation of sown grasslands, with high content of perennial legumes on meadows and arable lands, which are derived from intensive cultivation is one of the most promising areas of organic onion growing. Productivity of single-species crops of perennial legumes and their mixtures with boneless stalk on average for five years after the yield of 1 ha of dry mass in the version without fertilizers ranged from 5.12 to 8.38 tons, which is 1.7–2.5 times more in comparison with a grassy grass stand which is formed from one-species crops of a stokolos of a boneless. The highest productivity and level of symbiotic nitrogen accumulation (191–266 kg / ha) were provided by agroecosystems with the participation of alfalfa, and the lowest - meadow clover. The efficiency of soil liming during sowing of grasses and application of inoculation of seeds of perennial legumes with nitrogen-fixing and phosphorus-mobilizing preparations is given. Changes in the chemical composition of feed under the influence of the studied factors are also shown. **Conclusions.** Different types of perennial legumes and their mixtures with boneless stalk without fertilizers provide productivity in the range of 5.12–8.38 t / ha of dry weight, which is 1.7–2.5 times more than the grass cover, which is formed from single-species sowing of stokolos bezzostoy. Inoculation of perennial legume seeds with nitrogen-fixing and phosphorus-releasing microbiological preparations increases the productivity of perennial legumes and their mixtures with cereals by 2–6%, and soil liming before sowing – by 3–11%. The greatest effect is provided by the combined use of these bacterial drugs. Alfalfa sowing responds best to their use and liming.

Key words: perennial legumes, bacterial preparations, organic onion growing, productivity, dry mass, liming.

Introduction (problem statement). Due to the deepening ecological crisis, there is a steady trend of increasing demand for food organic production of agricultural products. Organic grassland farming is not only a set of measures aimed at obtaining environmentally safe forage. Organic grassland farming is closely integrated into the context of sustainable development and environmental management. It also includes a set of measures to preserve the environment, where a huge environmental and stabilizing role in agrolandscapes

meadows, protecting the soil from erosion and sources from siltation and pollution (Kurgak, 2013). It is the reduction of ploughed land, the reproduction of onion grasslands and a network of sanctuaries, nature reserves, hunting grounds, along with an increase in the production of organic fodder, helps to improve the ecological balance of the environment, the conservation of biodiversity of plants, animals, in particular species included in the Red Book (Kurgak, 2019; Mikhalchuk, 2010; Demyanchik, 2012). Taking into account the huge

unused potential of Ukrainian grasslands, (about 8 million hectares with migrations), in the direction of obtaining organic fodder products on them and transfer them from the category of the so-called «abandoned» lands in the fodder-producing, nature conservation, recreational and ecological zones. This will not only turn them into exemplary fodder crops and at the same time conservation elements of agrolandscapes on the basis of adaptive conservation ecological and recreational land use (Kurgak, 2019).

Natural forage lands in the vast majority as an ecologically clean area and a source of environmentally safe forage is an important component of ecological tourism, in combination with organic production meets the requirements of sustainable development and rational nature management (Parente, 2012; Kurgak, 2013). In line with the development of the tourism industry it is necessary to design and implement a program for the development of ecological tourism, and in particular agritourism, in which considerable attention is paid to the quality of food as an important element of the quality of service.

An integral component of organic production is the development and implementation of a set of measures aimed at the conservation of flora and fauna of the onion grasslands by creating a network of nature reserves, wildlife refuges, hunting and deer farms with keeping and grazing animals in cages (Tyshkevich, 2010). In organic fodder production great attention should be paid to quality control of fodder by certified laboratories. Fodder should meet the requirements of state standards of Ukraine, where control is provided not only on the main indicators of quality, but also on safety indicators (Kurgak, 2013, 2019; Demidass, 2020).

In recent years, due to the increasing processes of climate aridization and xerophytization of vegetation to ensure sustainable development there is a need to search and introduce drought-resistant species from the group of mesoxerophytes and xeromesophytes in the Forest-steppe and even Polesie for stable organic fodder production should be introduced drought-resistant species of fodder plants or irrigation (Petrychenko, 2012). Organic feed production combined with the corresponding direction of livestock, is an inseparable link between crop and livestock production and an important element of organic production in general. For Ukraine, the development of organic feed production should go hand in hand with the development of cattle breeding and livestock and increase the number of

cattle and not only the dairy, but the meat direction. Livestock provides an opportunity to obtain organic fertilizers, which are the source of receipt of nutrients in the soil for organic production. Optimal load per 1 hectare of forage land to bring up to 1.5 cattle or 15 sheep, which will produce and introduce 10 tons of organic matter into the soil, which will ensure high and sufficient yields not only crops forage group, but also other crops (Kurgak, 2013).

Natural forage lands and hayfields and pastures placed on them are a complex integral system, in which all components are functionally closely connected with each other by exchange processes of substance and energy flows. Laws, which are characteristic for natural meadows, are in the basis of existence and sowing meadows (Mirkin, 1986; Kurgak, 2010, 2017; Petrichenko, 2013). When forming sowing herbage, as a rule, perennial grasses and legumes should be included before herbage mixtures, taking into account the basic principles of component selection. Creation of sown grasses with an increased content of legumes is one of the most promising directions of organic grassland farming in the world (Hamacher, 2016; Weiß, 2016; Wallenhammar, 2018; Karbivska, 2020). Inclusion of legume grasses in legume-grass cenoses without mineral nitrogen application increases the productivity of grasslands by 1.5–2.5 and by 2–3 times in terms of protein collection compared to cereal grasses (Nilsdotter-Linde, 2016; Karbivska et al., 2020a, 2020b; Kovtun, 2020). It is known that legume grasses are not durable. Therefore, the main provisions and practical measures to increase their productive longevity in sown meadow legume-grass agrocenoses are formulated (Kurgak, 2010, 2018, 2020; Petrichenko, 2013; Damborg, 2016; Hynes, 2018). The productive longevity of legumes is influenced not only by the correct selection of components, but also by the ways of placement of cereal and legume components (the best way with alternate placement of cereal and legume components in separate rows from non-widespread rows or narrow strips), replacement of mineral and symbiotic nitrogen by years of use and the like.

Despite the significant amount of research issues of substantiation of conceptual and technological foundations of organic grassland in Ukraine and identifying the role of perennial legumes in increasing the productivity of meadow phytocenoses until recently remain insufficiently studied, which was the subject of our research, are covered in the article.

Materials and methods. Field studies were conducted during 2016–2020. In the National Scientific Center «Institute of Farming of the National Academy of Agrarian Sciences of Ukraine» (northern part of the Forest-steppe of Ukraine, Chabany settlement, Kiev-Svyatoshinsky district of Kiev region). Soil of the research site is the dark gray podzolized coarse-piluvate light loam on loess-like loam. The depth of humus horizon is 35–40 cm. Humus content in 0–20 cm layer was 2.4%, pH 5.2, hydrolytic acidity 4.2 mg-eq/100 g, lunohydrolyzed nitrogen 13.1, labile phosphorus 17.1, and exchangeable potassium 12.9 mg per 100 g of soil. Depth of groundwater was about 3 m. Relief was flat with a barely noticeable slope towards the Dnieper River.

The climate of the region where the study was conducted is moderately continental. Average annual air temperature is 6–7 °C. Average annual precipitation is 480–620 mm. Weather conditions in Chabany settlement, during the years of research (2016–2020) were favorable for growth and development of perennial grasses, despite the fact that during the year and the growing season air temperature exceeded by 1.5–2.0 °C, and the amount of precipitation was less than the average perennial rate.

Scheme of experiments is given in the following tables. Area of the sowing plot is 30 m, recording area is 25 m². Repetition of the experiment is four times. Cultivation technology of perennial grasses and their mixtures, excluding the studied factors was generally accepted. Nitrogen fertilizers in the form of ammonium nitrate with the active substance content of 34%, phosphorus as simple superphosphate (18.7%) and potassium as potassium chloride (60%), according to the experiment scheme, were applied annually in early spring. Liming was carried out by lime application under pre-sowing cultivation during perennial grass sowing in a dose of 5 t/ha. Seeds of perennial legume grasses were inoculated with strains of nodule bacteria, according to the technological instructions, immediately before sowing. Dry weight content was determined by thermostatic weight method at 105 °C. The yield of green mass was accounted by weight method, by weighing with subsequent recalculation of yield from 1 ha of dry weight (Dospekhov, 1985).

The purpose of research. To establish the best perennial legumes when grown in single-species and compatible fodder agrocenoses with stokolos boneless and the effectiveness of the use of nitrogen-fixing and

phosphormobilizing bacterial preparations, as well as liming.

Research results and their discussion. Analysis of the research results showed that in all research variants the best preserved grasses in five years were sown alfalfa and hornwort (Table 1). The average share of these plants when liming and simultaneous application of nitrogen-fixing and phosphate-mobilizing bacterial preparations in single-species crops was 80–84% and in mixtures with awnless brome grass – 65–78%. The average proportion of eastern brome grass was at 59% and 45%, respectively, which, unlike other legumes, was greatest in years 3–5 of use. The lowest share of the seeding crop was in agrocenoses with meadow clover (48 and 35%, respectively), which remained at a high level only in the first three years of vegetation, later it almost completely fell out of the grass stand. In our studies, the accumulated symbiotic nitrogen ranging from 95 to 266 kg/ha against the background of liming and joint application of nitrogen-fixing and phosphate-mobilizing bacterial preparations, different species of perennial legume grasses in single and joint crops on average for 2016–2020 (see Table 1). Legume grass species both in single-species crops and in mixtures with awnless brome grass accumulated more symbiotic nitrogen. Alfalfa in single-species crops accumulated the most nitrogen, while meadow clover in mixtures with awnless brome accumulated the least. All species of perennial leguminous grasses accumulated more of it in single-species crops than in mixtures with cereals. The productivity of single-species crops of perennial legumes and their mixtures with awnless brome grass with yield of dry mass per 1 ha in the variant without liming and without application of nitrogen fixing preparations, due to the action of symbiotic nitrogen, was high and ranged from 5.12–8.38 tons on average within 2012–2020 (Table 1). This productivity was 1.7–2.5 times higher in comparison with the grass herbage formed from single-species awnless brome grass crops.

The most productive was the agrophytocenosis that was formed on the basis of single-species alfalfa crops, and the least productive was the agrocenosis that was formed on the basis of a mixture of meadow clover and awnless brome grass. On average for five years legume-grass herbage in relation to equal-species legume crops, formed a total for three mowing by 3–9% less productivity from 1 hectare of dry weight. The grass stand formed from awnless brome grass formed the lowest

Table 1. Productivity of perennial leguminous grasses and their mixtures with awnless brome grass depending on soil liming and application of bacterial preparations, t/ha (average for 2016-2020rr).

Grass species and sowing rate, kg/ha	Liming	Legume share, %*	Nc**, kg/ha	Bacterial preparation productivity			
				None	nitrogen-fixing	phosphate-mobilizing	nitrogen-fixing and phosphate-mobilizing
Meadow clover – 20	–	48	132	5.12	5.24	5.22	5.30
	+			5.23	5.39	5.21	5.55
Alfalfa – 20	–	84	266	8.38	8.71	8.61	8.92
	+			9.27	9.73	9.36	9.91
birdsfoot deer vetch– 14	–	89	187	6.95	7.16	7.15	7.18
	+			7.10	7.27	7.24	7.43
Eastern galega – 30	–	59	177	6.27	6.56	6.33	6.70
	+			6.92	7.08	6.89	7.37
Meadow clover –10 + Anwless brome grass – 15	–	35	95	5.37	5.53	5.40	5.57
	+			5.42	5.67	5.47	5.84
Alfalfa – 10 + Anwless brome grass –15	–	65	191	7.73	8.13	7.98	8.18
	+			8.61	8.82	8.69	8.95
birdsfoot deer vetch –7 + Anwless brome grass –15	–	78	140	6.69	7.02	6.85	7.02
	+			7.05	7.14	7.01	7.33
Eastern galega –15 + Anwless brome grass – 15	–	45	135	6.03	6.16	6.06	6.32
	+			6.54	6.68	6.57	6.89
Anwless brome grass – 30 + N ₁₂₀	–	–	–	5.60	–	–	–
Anwless brome grass – 30 (control)	+	–	–	2.81	–	–	–
HIP ₀₅ , t/hadry weight: standing grass – 0.36, bacterial preparation – 0.38, liming – 0.32. <i>*Share of legume component and accumulation of symbiotic nitrogen by leguminous grasses (Nc**) are given on the background of liming and joint application of nitrogen-fixing and phosphate-mobilizing bacterial preparations.</i>							

productivity of dry mass (2.81 t/ha) without fertilizers. After application of N120 the productivity of awnless brome grass increased 2.2 times and was 5.60 t/ha of dry mass.

The research revealed a positive effect on the productivity of perennial legumes and their mixtures with awnless brome grass by inoculating the seeds with nitrogen-fixing and phosphorus-mobilizing bacterial strains during sowing. Although the effect of bacterial preparations on the development of legume species weakened

somewhat over the years, the productivity of perennial legume grasses and legume-straw grasses increased by an average of 2-6% from their action. Alfalfa responded better to the application of bacterial preparations. Its productivity on joint application of nitrogen-fixing and phosphate-mobilizing bacterial preparations was the highest of all the studied variants and reached 9.91 t/ha of dry weight. Somewhat greater productivity gains from the use of bacterial preparations were on the background of liming.

Studies also revealed a positive effect of liming the soil on the productivity of perennial legumes and their mixtures with awnless brome. Against the background of liming all legumes and legume-grasses provided 3–11% higher productivity than the cenoses of these grasses without liming. Both in single-species and joint crops with awnless brome responded better to lime treatment. Legume-grass grasses over the years after reducing the legume component in the grasses reacted worse to lime treatment. Legume-grass herbage grasses responded to lime treatment somewhat weaker than pure legume crops.

Productivity of legume and legume-grass stands changed significantly depending on the time of existence and use of grasses (Table 2). Liming had a positive effect on the productivity of agrocenoses in all years. In

all years, single-species crops of perennial leguminous grasses were more productive than their joint crops with awnless brome. The exceptions were agrocenoses that were formed with meadow clover, whose joint crops with awnless brome starting from the third year were more productive than grasses that were formed on the basis of single-species clover crops.

The lowest productivity was in 2016. In this year, grass crops for spring barley ranged from 0.9–2.66 t/ha of dry weight. The least responsive to the presence of the cover crop in the year of sowing were hornbeam and meadow clover, which provided the highest fodder productivity after mowing the cover crop, which was at 2.32–2.66 t/ha dry weight. In the year of sowing, the cover crop negatively affected the productivity of agrocenoses when adding awnless brome to perennial legumes, which

Table 2. Productivity of perennial leguminous grasses in single-species crops and mixtures with awnless brome by years of research depending on liming, t/ha dry weight*

Grass specie and sowing rate, kg/ha	Liming	Year				
		2016	2017	2018	2019	2020
Meadow clover – 20	–	2.32	9.37	8.27	4.04	2.50
	+	2.47	10.02	8.62	4.10	2.54
Alfalfa – 20	–	2.14	8.58	12.75	11.35	9.8
	+	2.25	10.30	13.43	12.63	10.89
Birdsfoot deer vetch – 14	–	2.54	9.02	9.82	8.30	6.22
	+	2.66	9.12	10.03	8.83	6.51
Eastern galega – 30	–	1.02	6.41	9.31	8.51	8.28
	+	1.09	6.79	9.89	9.77	9.31
Meadow clover – 10 + Anwless brome grass – 15	–	1.51	7.59	7.92	7.72	3.12
	+	1.52	8.05	8.48	7.88	3.28
Alfalfa – 10 + Anwless brome grass – 15	–	1.46	7.63	11.63	10.83	9.38
	+	1.89	8.51	11.97	11.84	10.55
Birdsfoot deer vetch – 7 + Anwless brome grass – 15	–	1.66	8.56	9.62	8.07	7.19
	+	1.92	8.63	9.73	8.48	7.89
Eastern galega – 15 + Anwless brome grass – 15	–	0.91	6.47	8.47	7.6	8.15
	+	1.21	6.71	8.65	8.32	9.57
HIP ₀₅ , t/ha						
Stand grass		0.21	0.39	0.43	0.41	0.37
Liming		0.20	0.32	0.41	0.35	0.33

**Productivity indicators are given against the background of joint application of nitrogen-fixing and phosphate-mobilizing bacterial preparations.*

decreased in comparison with single-seeded crops. In 2017, the second year of crop life, the productivity of single-species crops of perennial legume grasses and their mixtures with cereals ranged from 6.41–10.30 t/ha of dry weight. This year, agrocnoses with meadow clover, hornwort, and alfalfa were the most productive, both in single-species crops and in mixtures with awnless brome grass. The least productive were the agrocnoses which were formed with eastern goatweed.

The highest level of productivity was provided by legumes and legume-grass agrocnoses in 2018. – in the third year of their use, in the range of 7.92–13.43 t/ha of dry weight. This year in terms of productivity, the difference between different herbage stands with the participation of different types of legume grasses was the smallest. This is due to the fact that along with the good preservation in the herbage of alfalfa and hornwort, meadow clover was well preserved in the

agrocnoses, and also powerfully grew and increased its share in the yield of goatweed. The best response to liming, as well as to the application of bacterial preparations this year was lucerne sowing, when its productivity under the influence of these factors increased by 1.1-1.2 times and reached a maximum value of 13.43 t/ha of dry weight at the combined application of lime, nitrogen-fixing and phosphate-mobilizing preparations.

In 2019, the fourth year of vegetation, the productivity of legume and legume-grass agrocnoses ranged from 4.00–12.63 t/ha dry weight. Both between single-species crops and joint crops of legumes and awnless brome grass, the highest productivity was provided by agrocnoses with alfalfa. Productivity of agrocnoses formed on the basis of meadow clover, due to its loss from herbage was low and approached the productivity of cereal stoloniferous grass. Productivity of Oriental goatweed approached to that of hornwort.

Table 3. Chemical composition and digestibility of forage mass by grass depending on the botanical and species composition of the cenosis, % (average for 2016–2020)

Grass specie and sowing rate, kg/ha	Liming	Crude Protein	Protein	Crude Fat	Crude Metal	NFE	P ₂ O ₅	K ₂ O	Digestibility
Meadow clover – 20	–	19,6	18,3	2,9	24,7	45,2	0,82	2,84	66
	+	20,7	18,6	3,1	24,1	45,4	0,83	2,93	64
Alfalfa – 20	–	19,5	18,3	3,2	24,7	44,5	0,80	3,02	72
	+	19,9	18,2	3,1	23,1	46,0	0,84	3,48	70
birdsfoot deer vetch – 14	–	19,3	17,9	3,2	23,3	45,7	0,85	3,17	73
	+	20,2	18,4	3,3	24,4	44,6	0,87	3,08	71
Eastern galega – 30	–	18,8	17,0	3,1	24,8	45,6	0,87	2,83	68
	+	19,2	17,5	3,4	24,7	45,8	0,88	2,89	73
Meadow clover – 10 + Anwless brome grass – 15	–	15,8	14,4	2,6	25,3	48,6	0,79	2,93	67
	+	15,7	14,2	2,9	25,1	47,9	0,81	2,84	66
Alfalfa – 10 + Anwless brome grass – 15	–	16,7	15,3	2,7	24,9	48,1	0,83	2,95	64
	+	17,0	15,5	3,0	25,4	47,1	0,81	3,01	68
birdsfoot deer vetch – 7 + Anwless brome grass – 15	–	16,8	15,3	2,9	24,7	47,9	0,85	2,81	67
	+	16,5	15,5	2,9	25,6	47,4	0,83	2,98	63
Eastern galega – 15 + Anwless brome grass – 15	–	15,9	14,8	2,8	25,1	48,0	0,89	2,88	61
	+	16,8	15,0	2,8	25,5	47,5	0,91	2,80	64
Anwless brome grass – 30 (control)	–	11,4	11,7	2,4	26,0	49,8	0,71	2,54	58
Anwless brome grass – 30 + N ₁₂₀	–	14,8	12,3	2,5	26,2	49,2	0,82	2,68	60

Alfalfa responded best to lime treatment both in single-species crops and joint crops with awnless brome. Legume-grass herbage grasses in comparison with single-species legume crops formed by 5–7% lower productivity. The exception was meadow clover crops whose productivity, after its exclusion from the herbage was 1.8–2 times higher than in single-species crops.

A similar pattern was observed in 2020. This year compared with 2019, there was a tendency to decrease the productivity of these agrocenoses, excluding goat-grass and goat-grass-grass stands, the productivity of which did not change much. The greatest decrease in productivity occurred in phytocenoses, which were formed on the basis of straw-calf mixture. The productivity of swaths in the studied agrocenoses changed significantly. In all swaths high productivity was noted in agrophytocenoses which were formed on single-species alfalfa sowings. In all studied variants it was the highest in the first swath with parameters of 3.20–5.55 t/ha of dry weight, and the lowest in the third swath. In all swaths, the highest productivity was provided by joint application of nitrogen-fixing and phosphate-mobilizing bacterial preparations.

The chemical composition of the feed of perennial legumes and leguminous grasses on average over the years of research is given in table. 3. Analysis

The results showed that the presence of the legume component in the grassland, due to the action of symbiotic nitrogen, improved the chemical composition. In particular, in the variants without nitrogen fertilizers in the dry mass of feed, the content of crude protein increased from 11.4%, which was in the stock of grass, to 15.3–19.4% or 3.9–8.0%. At the same time, the content, a component of crude protein, protein increased by 2.4–5.8%, crude fat - by 0.2–0.5%, the mineral composition of feed and digestibility of dry matter in vitro improved. Thus, the content of P_2O_5 increased by 0.9–0.14%, K_2O – by 0.25–0.56%, and the digestibility of dry mass – by 2–12%. At the same time, the content of nitrogen-free extractives decreased by 1.3–4.2%.

The chemical composition of the dry mass of fodder was noticeably better in single-species crops of perennial legumes than in agrocenoses formed on the basis of legume-cereal mixtures.

A similar pattern with the chemical composition of the feed was under the action of mineral nitrogen fertilizers on cereals. In this case, the application of N120 in comparison with the option without its application

(without fertilizers), the content of crude protein increased by 3.4%, and protein – by 0.6%.

Analysis of the results of research on the chemical composition of feed shows that the best indicators of feed quality in terms of crude protein, protein, crude fat and digestibility of dry matter in vitro and mineral composition were characterized by single crops of alfalfa and lovable, as well as coenoses of clover.

During entire research period, we found that legume components in the legume-grass cenoses, depending on soil and climatic factors in the Polissya and the northern part of the steppe accumulate aboveground mass of the symbiotic fixed nitrogen in the range from 50 to 200 kg/ha. Average weighted values of its accumulation by leguminous grasses are shown in Table 4. Alfalfa and sandy sainfoin accumulate it most in joint crops with cereal species. Single-species crops of perennial legume grasses accumulate 20–30% more symbiotic nitrogen than legume-grass agrocenoses.

To enhance the ecological role of grasslands, surface improvement measures of herbaceous ecosystems are applied in combination with protection measures, which are to preserve a wide range of biodiversity of natural and sown grass vegetation, to protect soils from erosion, and water sources from pollution and siltation. It should be borne in mind that as a conservation measure on natural forage lands, it is desirable to apply phytoreclamation, when instead of chemical

Table 4. Approximate amount of symbiotic fixed nitrogen accumulated by legumes in meadow legume-grass phytocenoses, kg/ha

Grass	Symbiotic nitrogen
Sowing stand grass	
Alfalfa-grass	150–170
Meadow clover-grass	130–150
White clover-grass	110–130
Crowfoot-grass	100–120
Holy clover-grass	150–170
Natural stand grass	
Legume grass with various legumes	70–100

Data are given for legume content of (50-60)%. At legume content of 30-50% the symbiotic nitrogen is accumulated 1.2-1.5 times less, and at 60-80% - 1.2-1.5 times more. Data are given for optimal soil and climatic conditions for a particular type of legume component. Sown alfalfa, hornwort, and sandy sainfoin were used in the experiments.

ameliorants (lime, dolomite flour, and gypsum) which pollute the environment, perennial grasses resistant to increased acidity or soil salinity are used. Phytomeliorative ability on saline soils is characterized by melilot, which, in addition to high yields, removes toxic salts.

In organic production on forage lands it is advisable to apply organic fertilizers (manure, green manure, etc.) in doses of 10–30 t/ha in the amount of manure in light texture soils. The dose of application of litterless manure (semi-liquid, liquid and manure runoff) is calculated by the content of nitrogen or nutrient element (nitrogen, phosphorus or potassium), which is at its maximum (Kurgak, 2010).

CONCLUSIONS

The conceptual and technological basis of organic grassland management provides a set of measures for the production of environmentally safe forage for animals with the maximum use of biological factors of intensification and aimed at the conservation of the environment, in particular soils, water sources and biodiversity. Creation of sown grasses with an increased content of perennial leguminous grasses is one of the most promising areas of organic grassland farming.

We have established that the productivity of monoecious leguminous grasses and their mixtures with awnless brome grass was high and varied within the range of 5.12–8.38 tons during five years on the average per

1 ha of dry weight in the variant without fertilization, which is 1.7–2.5 times higher than the productivity of grass stand formed from monoecious grasses with awnless brome grass. The highest productivity and the level of accumulation of symbiotic nitrogen (191–266 kg/ha) are provided by agrocenoses with lucerne sowing and the lowest one with meadow clover. Inoculation of perennial legume grass seeds with nitrogen-fixing and phosphate-mobilizing preparations increases the productivity of perennial legume grasses and their mixtures with cereals by 2–6%. The greatest effect is provided by joint application of these preparations. Soil liming before sowing perennial leguminous grasses and their mixtures with cereals increases their productivity by 3–11%. Alfalfa responds best to liming and the use of bacterial preparations.

The presence of legumes in the grass, due to the action of symbiotic nitrogen, improves the chemical composition of the feed. In particular, in the variants without the application of nitrogen fertilizers in the dry weight of the feed, the crude protein content increases from 11.4% to 15.3–19.4% or by 3.9–8.0%. The protein content increases by 2.4–5.8%, crude fat - by 0.2–0.5%, improves the mineral composition of feed and digestibility of dry matter *in vitro*.

LITERATURE

1. Damborg V.K., Stødkilde L., Jensen S.K. and Weisbjerg M.R. Characterisation of protein and fibre in pulp after biorefining of red clover and perennial ryegrass. The multiple roles of grassland in the European bioeconomy: Proceedings of the 26th General Meeting of the European Grassland Federation. Trondheim, Norway. 4-8 September. 2016. P. 366–371.
2. Демидась Г.І., Галушко І.В. Мінеральний склад кормової маси різних сортів конюшини лучної залежно від елементів технології вирощування. Корми і кормовиробництво. № 89. 2020. С.151–160. <https://doi.org/10.31073/kormovyrobnytstvo202089-15>.
3. Демянчик В.Т. Динамика экологического и фаунистического потенциала мелиоративных систем в режиме подтопления. Природное асыроддзэ Палесся: асаблівасці і перспектывы развіцця. Зб. н. п. Палескі аграрна-экалагічны інстытут НАНБ. Брэст: Альтернатива. Вып. 5. 2012 С. 74–78.
4. Доспехов Б. А. Методика полевого опыта. Москва. Агропромиздат. 1985. 347 с.
5. Hynes D., Laidlaw S. and Ferris C. Effect of management strategy on wilting of monocultures and mixture of red clover and perennial ryegrass // Sustainable meat and milk production from grasslands: Proceedings of the 27th General Meeting of the European Grassland Federation. Cork, Ireland. 17–21 June, 2018. P. 249–251.
6. Карбівська У.М. Накопичення кореневої маси бобових трав та її вплив на поживний режим дерново-підзолистого ґрунту. Передгірне та гірське землеробство і тваринництво. Вип. 67 (I). 2020 С. 84–97. [https://www.doi.org/10.32636/01308521.2020-\(67\)-1-6](https://www.doi.org/10.32636/01308521.2020-(67)-1-6).
7. Karbivska U., Kurhak V., Gamayunova V., Butenko A. et. al. Productivity and quality of diverse ripe

- cereal grass fodder depending on the methods of soil cultivation. *Acta Agrobotanica* (Vol. 73. №3. 2020. P. 1–11.) Article ID: 7334 DOI: 10.5586/aa.7334.
8. Karbivska, U.M., Kurgak, V.G., Kaminskyi, V.F., Butenko, A.O., Davydenko, G.A., Viunenko, O.B., Vyhanaiilo, S.M., Khomenko, S.V. Economic and Energy Efficiency of Forming And Using Legume-Cereal Grass Stands Depending on Fertilizers. *Ukrainian Journal of Ecology*, 10 (2). 2020. 284–288. DOI: 10.15421/2020_98.
9. Ковтун К.П., Векленко Ю.А., Ящук В.А. Формування фітоценозу та продуктивності еспарцето-злакових травосумішок залежно від способів сівби та просторового розміщення видів в умовах Лісостепу правобережного. *Корми і кормовиробництво*. № 89. 2020. С.112–120. <https://doi.org/10.31073/kormovyrobnytstvo202089-11>.
10. Кургак В.Г. Лучні агрофітоценози. Київ: ДІА. 2010. 374 с.
11. Кургак В.Г., Кулаковская Т.В. Использование лугопастбищных угодий в органическом производстве Европы. Поєднання науки, освіти, практичного виробництва і реалізація якісної органічної продукції: мат. ІУ Міжн. наук.-практ. конф., 2013. С. 156–168.
12. Кургак В.Г. Використання багаторічних бобових трав на луках України за органічного виробництва трав'яних кормів. Поєднання науки, освіти, практичного виробництва і реалізація якісної органічної продукції: Матеріали міжнародної науково-практичної конференції 26 червня 2013 р. Київ-Іллінці. С. 169–178.
13. Кургак В. Г., Волошин В. М. Підвищення ефективності використання багаторічних бобових трав на луках України. Посібник українського хлібороба «Біологізація землеробства»: науково-практичний збірник. Київ. Том 1. 2017. С. 288–291.
14. Кургак В.Г. Способи підвищення продуктивного довголіття бобово-злакових лучних агрофітоценозів. Поєднання науки, освіти, практичного виробництва і реалізація якісної органічної продукції : м-ли 9 міжнародної науково-практичної конференції 2018 р. Київ-Іллінці. 2018. С 9–17.
15. Кургак В.Г., Карбівська У.М. Особливості формування бобово-злакових агрофітоценозів на дерново-підзолистих ґрунтах Прикарпаття України. *Корми і кормовиробництво*. 2020. № 89. С. 121–133. <https://doi.org/10.31073/kormovyrobnytstvo202089-12>
16. Миркин Б.М. Что такое растительные сообщества. Москва: Наука. 1986. 164 с.
17. Михальчук Н.В., Галуц О.А., Ковалев И.В., Мороз В.А., Савчук С.С. Жемчужины природы Малоритчины. Брест: Альтернатива, 2010. 40 с.
18. Nilsdotter-Linde N., Halling M.A. and Jansson J. Widening the harvest window with contrasting grass-clover mixtures. The multiple roles of grassland in the European bioeconomy: Proceedings of the 26th General Meeting of the European Grassland Federation. Trondheim, Norway. 4–8 September 2016. P. 191–193.
19. Petrychenko V., Bohovin A. and Kurhak V. More efficient use of grassland under climate warming. Grassland – a European Resource?: Proceedings of the 24th General Meeting of the European Grassland Federation. Lublin, Poland. 3–7 June 2012. P. 151–153.
20. Петриченко В.Ф., Кургак В.Г. Культурні сіножаті та пасовища України. Київ: Аграрна наука. 2013. 432 с.
21. Parente G., Bovolenta S. The role of grassland in rural tourism and recreation in Europe . In: Golinski P., Warda M. and Stypinski P. (ed.) Grassland – a European Resource. Proceedings of the 24th EGF General Meeting. Lublin (PL) 3–7 June 2012. Volume 17. Grassland Science in Europe. P. 733–743.
22. Тышкевич В.Е. Восстановление агроценозов и луговое хозяйство на торфяниках для интенсификации охотничьего хозяйства Беларуси. Мелиорация. 2010. № 2 (64). С. 178–191.
23. Wallenhammar A., Stoltz E., Omer Z. and Granstedt A. Production capacity of forage legumes and persistence to root rot in organic mixed swards. Sustainable meat and milk production from grasslands: Proceedings of the 27th General Meeting of the European Grassland Federation. Cork, Ireland. 17–21 June, 2018. P. 375–377.
24. Weiß K. and Kalzendorf C. Effect of wilting and silage additives on silage quality of lucerne, red clover and legume-grass mixtures . The multiple roles of grassland in the European bioeconomy. Proceedings of the 26th General Meeting of the European Grassland Federation. Trondheim, Norway. 4–8 September 2016. P. 170–172.

REFERENCES

1. Damborg, V.K., Stødtkilde, L., Jensen, S.K., Weisbjerg, M.R. (2016). Characterisation of protein and fibre in pulp after biorefining of red clover and perennial ryegrass. The multiple roles of grassland in the European bioeconomy: Proceed. XXVI General Meeting of the European Grassland Federation. Trondheim, Norway. 4-8 September, 366–371 [in English].
2. Demidas, G.I., Galushko, I.V. (2020). Mineral composition of fodder mass of different varieties of meadow clover depending on the elements of cultivation technology. *Feed and feed production*, 89, 151–160. <https://doi.org/10.31073/kormovyrobnytstvo202089-15> [in Russian].
3. Demyanchik, V.T. (2012). Dynamics of ecological and faunal potential of reclamation systems in the mode of flooding. Natural environment of Polesie: features and prospects of development. Coll. N. Polesie Agrarian and Ecological Institute of the National Academy of Sciences of Belarus, 5, 74–78 [in Russian].
4. Dospekhov, B.A. (1985). Methods of field experience. Moscow. Agropromizdat. Hamacher M., Loges R. and Taube F. (2016). Evaluation of fifteen leguminous and non-leguminous forage species to improve forage quality of temporary grasslands in northern Germany. The multiple roles of grassland in the European bioeconomy: Proceed. XXVI General Meeting of the European Grassland Federation. Trondheim, Norway. 4–8 September, 263–265 [in Russian].
5. Hynes, D., Laidlaw, S., Ferris, C. (2018). Effect of management strategy on wilting of monocultures and mixture of red clover and perennial ryegrass. Sustainable meat and milk production from grasslands: Proceed. XXVII General Meeting of the European Grassland Federation. Cork, Ireland, 17–21 June 249–251 [in English].
6. Karbivska, U.M. (2020). Accumulation of root mass of legumes and its influence on the nutrient regime of sod-podzolic soil. *Foothill and mountain agriculture and animal husbandry*, 67(1), 84–97. [https://www.doi.org/10.32636/01308521.2020-\(67\)-1-6](https://www.doi.org/10.32636/01308521.2020-(67)-1-6) [in Ukrainian].
7. Karbivska, U., Kurhak, V., Gamayunova, V., Butenko, A. (2020a) Productivity and quality of diverse rye-cereal grass fodder depending on the methods of soil cultivation. *Acta Agrobotanica*, 73(3), 1–11. DOI: 10.5586/aa.7334 [in Ukrainian].
8. Karbivska, U.M., Kurgak, V.G., Kaminskyi, V.F., Butenko, A.O., Davydenko, G.A., Viunenko, O.B., Vyhanaiilo, S.M., Khomenko, S.V. (2020b). Economic and energy efficiency of forming and using legume-cereal grass stands depending on fertilizers. *Ukrainian Journal of Ecology*, 10 (2), 284–288. DOI: 10.15421/2020_98 [in Ukrainian].
9. Kovtun, K.P., Veklenko, Yu.A., Yashchuk, V.A. (2020). Formation of phytocenosis and productivity of sainfoin-cereal grass mixtures depending on the methods of sowing and spatial distribution of species in the conditions of the right-bank Forest-Steppe. *Feed and feed production*, 89, 112–120. <https://doi.org/10.31073/kormovyrobnytstvo202089-11> [in Ukrainian].
10. Kurgak, V.G. (2010). Meadow agrophytocenoses. Kyiv: DIA [in Ukrainian].
11. Kurgak, V.G. (2013). The use of perennial legumes in the meadows of Ukraine in the organic production of fodder. Combination of science, education, practical production and sale of quality organic products. Proceed. Int. Sc. Conf, June 26. Kyiv-Illintsi, 169–178 [in Ukrainian].
12. Kurgak, V.G., Karbivska, U.M. (2020). Peculiarities of formation of leguminous-cereal agrophytocenoses on sod-podzolic soils of Prykarpattia of Ukraine. *Feed and feed production*, 89, 121–133. <https://doi.org/10.31073/kormovyrobnytstvo202089-12> [in Ukrainian].
13. Kurgak, V.G., Kulakovskaya, T.V. (2013). Use of meadow pastures in organic production in Europe. Combination of science, education, practical production and sale of quality organic products. Matt. Yiwu Int. Sc. Conf., 156–168 [in Ukrainian].
14. Kurgak, V.G., Voloshin, V.M. (2017). Improving the efficiency of perennial legumes in the meadows of Ukraine. Handbook of Ukrainian farmers «Biologization of agriculture». Proceed. Sc. Conf. Kyiv, 1, 288–291 [in Ukrainian].
15. Kurgak, V.G. (2018). Ways to increase the productive longevity of legume-cereal meadow agrophytocenoses. Combination of science, education, practical production and sale of quality organic products. Proceed. IX Int. Sc. Conf., Kyiv-Illintsi, 9–17 [in Ukrainian].
16. Mikhalchuk, N.V., Galuts, O.A., Kovalev, I.V., Moroz, V.A., Savchuk, S.S. (2010). Pearls of nature Maloritchina. Brest: Alternativa [in Russian].

17. Mirkin, B.M. (1986). What are plant communities. Moscow. Nauka [in Russian].
18. Nilsdotter-Linde, N., Halling, M.A., Jansson, J. (2016). Widening the harvest window with contrasting grass-clover mixtures. The multiple roles of grassland in the European bioeconomy: Proceed. XXVI General Meeting of the European Grassland Federation. Trondheim, Norway. 4-8 September, 191–193[in English].
19. Parente, G., Bovolenta, S. (2012). The role of grassland in rural tourism and recreation in Europe. In: Golinski P., Warda M. and Stypinski P. (ed.) Grassland – a European Resource. Proceed. XXIV EGF General Meeting. Lublin (PL) 3-7 June, *Grassland Science in Europe*, 17, 733–743[in English].
20. Petrichenko, V.F., Kurgak, V.G. (2013). Cultural hayfields and pastures of Ukraine. Kyiv: Agrarian Science [in Ukrainian].
21. Petrychenko, V., Bohovin, A., Kurhak, V. (2012). More efficient use of grassland under climate warming. Grassland – a European Resource? Proceed. XXIV General Meeting of the European Grassland Federation. Lublin, Poland. 3–7 June, 151–153 [in English].
22. Tyshkevich, V.E. (2010). Restoration of agrocenoses and meadow farming on peat bogs for intensification of hunting economy of Belarus. *Melioration*, 2 (64), 178–191[in Russian].
23. Wallenhammar, A., Stoltz, E., Omer, Z., Granstedt, A. (2018). Production capacity of forage legumes and persistence to root rot in organic mixed swards. Sustainable meat and milk production from grasslands: Proceed. XXVII General Meeting of the European Grassland Federation. Cork, Ireland. 17-21 June, 375–377 [in English].
24. Weiß, K., Kalzendorf, C. (2016). Effect of wilted grass silage additives on silage quality of lucerne, red clover and legume-grass mixtures. The multiple roles of grassland in the European bioeconomy. Proceed. XXVI General Meeting of the European Grassland Federation. Trondheim, Norway. 4–8 September, 170–172 [in English].

Кургак В.Г., Панасюк С.С.

Продуктивність багаторічних бобових трав в одновидових і сумісних із стоколосом безостим кормових агрофітоценозах

Мета. Встановити кращі багаторічні бобові трави при вирощуванні в одновидових і сумісних кормових агроценозах із стоколосом безостим та ефективність застосування на цих травостоях азотфіксувальних і фосформобілізувальних бактеріальних препаратів, а також вапнування. **Методи.** Загальнонаукові – гіпотез, індукції і дедукції, аналогії, узагальнення та спеціальні – польовий, лабораторний, математико-статистичний, розрахунково-порівняльний. **Результати.** На підставі узагальнення літературних джерел доведено, що створення сіяних травостой, із підвищеним вмістом багаторічних бобових трав на лучних угіддях та орних землях, які виводяться із інтенсивного обробітку є одним із найперспективніших напрямів ведення органічного лукувництва. Продуктивність одновидових посівів багаторічних бобових трав та їх сумішей зі стоколосом безостим в середньому за п'ять років за виходом з 1 га сухої маси у варіанті без внесення добрив коливалася у межах 5,12–8,38 т, що в 1,7–2,5 раза більше порівняно з злаковим травостоем, який сформовано із одновидового посіву стоколосу безостого. Найбільшу продуктивність та рівень нагромадження симбіотичного азоту (191–266 кг/га) забезпечили агроценози за участі люцерни посівної, а найменшу – конюшини лучної. Наведено ефективність вапнування ґрунту при сівбі трав та застосування інокуляції насіння багаторічних бобових трав азотфіксувальними та фосформобілізувальними препаратами. Показано також зміни хімічного складу корму під дією досліджуваних факторів. **Висновки.** Різні види багаторічних бобових трав та їх суміші зі стоколосом безостим без внесення добрив забезпечують продуктивність у межах 5,12–8,38 т/га сухої маси, що в 1,7–2,5 раза більше порівняно зі злаковим травостоем, який сформовано із одновидового посіву стоколосу безостого. Інокуляції насіння багаторічних бобових трав азотфіксувальними та фосформобілізувальними мікробіологічними препаратами підвищує продуктивність багаторічних бобових трав та їх сумішей із злаками на 2–6 %, а вапнування ґрунту перед сівбою – на 3–11 %. Найбільший ефект забезпечує поєднане застосування зазначених бактеріальних препаратів. Найкраще на їх застосування та вапнування реагує люцерна посівна.

Ключові слова: багаторічні бобові трави, бактеріальні препарати, органічне лукувництво, продуктивність, суха маса, вапнування.

ВІДОМОСТІ ПРО АВТОРІВ

Кургак В.Г., доктор с.-г. наук, професор, головний науковий співробітник ННЦ «Інституту землеробства НААН», вул. Машинобудівників, 2-б, смт Чабани, Києво-Святошинський р-н, Київська обл., 08162, e-mail: kurgak_luki@ukr.net, ORCID ID: <https://orcid.org/0000-0003-2309-0128>

Kurgak V.G., doctor of agricultural sciences Sciences, Professor, Chief Researcher of NSC “Institute of Agriculture NAAS”, st. Machine Builders, 2-b, village Chabany, Kyiv-Sviatoshynskyi district, Kyiv region, 08162, e-mail: kurgak_luki@ukr.net, ORCID ID: <https://orcid.org/0000-0003-2309-0128>

Панасюк С.С., кандидат с.-г. наук, старший науковий співробітник, старший науковий співробітник ННЦ «Інститут землеробства НААН», вул. Машинобудівників, 2-б, смт Чабани, Києво-Святошинський р-н, Київська обл., 08162, e-mail: petrivkas@ukr.net, ORCID ID: <https://orcid.org/0000-0003-4137-1169>

Panasyuk S.S., candidate of agricultural sciences Sciences, Senior Researcher, Senior Researcher NSC “Institute of Agriculture NAAS”, st. Machine Builders, 2-b, village Chabany, Kyiv-Sviatoshynskyi district, Kyiv region, 08162, e-mail: petrivkas@ukr.net, ORCID ID: <https://orcid.org/0000-0003-4137-1169>

Надійшла 16.06.2021